

IDENTIFICATION AND CHARACTERIZATION OF RUBBER DEGRADING ACTINOBACTERIA

H.M. RIFAAT*¹, M. A. YOSERY²

*e-mail: halamohamed6@hotmail.com

¹ *Microbial Chemistry Dept., National Research Center, Dokki, Cairo, Egypt*

² *Microbial Biotechnology Dept., National Research Centre, Dokki, Cairo, Egypt*

(Received 10th Nov 2003; accepted 12th May 2004)

Abstract. Of a great concern, the huge amount of waste rubber materials can cause environmental problems. Various methods have been proposed to solve this problem. One of those is the biodegradation of it by microorganisms. Bacteria able to degrade and use natural rubber latex as the sole source of carbon and energy were isolated from different ecosystems. 42 rubber-degrading bacteria were isolated. Out of these isolates, 31 were identified as *Streptomyces*, 5 as *Micromonospora*, 3 as *Actinoplanes*, 2 as *Gordonia* and 1 as *Nocardia* species. All rubber-degrading isolates were identified as members of Actinobacteria which is a large group of mycelium forming Gram-positive bacteria. Interestingly no Gram-negative bacteria could be isolated.

Keywords: *Streptomyces, Micromonospora, rubber degradation, isolation*

Introduction

Waste rubber is becoming a world wide waste disposal problem [13]. One particular concern is used it because of the huge number of natural rubber product produced and discarded annually and of potential environmental hazard should a NR stock pile catch on fire [14]. Consequently, it is very important and worth trying to develop a microbial process for waste NR disposal [6]. Natural rubber is consisting mainly of cis-1,4-polyisoprene and it is synthesized by more than 2000 plant species belonging mostly to the Euphorbiaceae. NR is still produced in large amounts (~ 10⁷ tons / year) from the rubber tree *Hevea brasiliensis*. The cis-1,4-polyisoprene, with an average molecular mass about 10⁶ Da, is the main constituent (> 90% of dry weight) of NR. NR is relatively resistant to microbial decomposition compared with many other natural polymers. Since the study by Sohngen and Fol, many reports have been published on the biodegradation of natural rubber by microorganisms [15, 10, 19, 18, 12 and 20]. NR contains a minimum of 90% rubber hydrocarbon together with small amounts of proteins, resins, fatty acids, sugars, and minerals [23]. Organic impurities in the rubber can support microbial growth [2 and 22]. Although many studies have been issued during the last decades on microbial degradation of rubber, only little is known about the occurrence of NR-degrading bacteria. Actinomycetes were almost the only organisms able to considerably decompose NR and use the hydrocarbon as a sole source of carbon and energy [4].

The present study was initiated to isolate and characterize a number of NR-degrading bacteria from various ecosystems in Egypt. It also suggests that rubber-degrading bacteria might be useful for the disposal of discarded rubber products. Identification and development of rubber metabolizing microorganisms potentially could provide a biotechnological solution to this problem.

Materials and Methods

Sampling sites

This study has concentrated on isolation of NR degrading actinobacteria from some localities in Egypt (Fig 1). The sampling sites were collected from various ecosystems (soils and fresh water from the River Nile as well as its bottom sediments).




Figure 1. Location map

Natural rubber source

Latex of *Hevea brasiliensis* was obtained from Weber and Schaer (Hamburg, Germany).

Isolation and Identification of NR-degrading bacteria

Microorganisms were isolated on mineral salts medium {8.0 g K_2HPO_4 , 1.0 g KH_2PO_4 , 0.5 g $(NH_4)_2SO_4$, 0.2 g $MgSO_4 \cdot 7H_2O$, 0.1g NaCl, 0.1g $Ca(NO_3)_2$, 20 mg $CaCl_2 \cdot 2H_2O$, 20 mg of $FeSO_4 \cdot 7H_2O$, 0.5 mg $Na_2MoO_4 \cdot H_2O$ and 0.5 mg $MnSO_4$ / L of deionised water} containing 25 to 100 mg of yeast extract and 20 g of agar / L that had been surface coated with a thin film (20-30 mg) of natural rubber. A hexane solution was also applied at the same medium and it was allowed to evaporate under a microbiological hood. Samples from different localities of Egypt from various ecosystems (soils, fresh water and their bottom sediments) were collected. The samples were serially diluted with sterile mineral medium and spread onto mineral plates coated with rubber and hexane and incubated for several weeks at 28 °C. The observed colonies were streaked onto the same rubber-coated plates until pure cultures were obtained.

Taxonomic characterisation of isolates

The isolates were characterised according to [1, 21 and 17] schemes based on their macro- and micromorphological properties. Adequate phenotypical tests set and chemotaxonomical investigations were used for the identification of strains including colony and micromorphological characteristics, pigment production tests, whole cell sugar pattern, cell wall chemotype, lecithinase, lipolysis, proteolysis, hydrolysis of pectin, chitin,

hippurate, casein, esculin, gelatine, degradation of xanthine, elastin, arbutin, utilization of sucrose, m-inositol, mannitol, L-rhamnose, raffinose, d-ribose, salicin, glucose, arabinose, fructose, xylose, galactose, nitrate reduction and H₂S production. The SPSS for Window release 6.0 statistical software has been used for clustering of the isolates, similarity calculations were based on simple matching coefficient (S_{SM} ; [16]). The results obtained were further evaluated using the above mentioned different systematic and determinative bacteriological manuals.

Qualitative assay method

Sugars (glucose, fructose, arabinose, sucrose, xylose, inositol, mannitol, rhamnose and raffinose) were sterilised by membrane filtration. 5% of each sugar was supplemented to the same mineral medium. Development or absence of the clearing zone formation was recorded.

Results and discussion

Isolation of NR-degrading bacteria

Ten samples from different ecosystems were screened for the presence of NR-degrading bacteria. 42 NR- degrading bacteria were isolated. They were identified as indicated by (i) size of the colonies developed on solid medium with purified NR latex as the sole source of carbon and energy in comparison to control a plate without NR and (ii) the appearance of translucent halos around the colonies. Interestingly all NR-degrading isolates belonged to the Actinobacteria and no Gram negative bacteria were isolated. Such results are in accordance with those reported by [4 and 20]. The first example of a Gram-negative rubber degrading bacterium, a *Xanthomonas* species, was reported by [19]. Linos et al. reported a new Gram- negative bacterium species namely *Pseudomonas aeruginosa* for NR-degradation [11]. However, our results cannot exclude NR-degradation capabilities encoded by Gram-negative bacteria in general. Potential Gram-negative NR-degraders might just require additional growth factors or degrade NR by co-metabolism.

Characterisation of NR-degrading bacteria

The isolates were divided into two main groups based on their colony morphology: the polysporic “streptomycetes” and the monosporic or non sporulating “other actinobacteria”. The members of the first group (31 strains) the isolates produced a yellow to grey coloured aerial mycelium with rectusflexiblis or spiral spore chains. The streptomycete isolates were identified as *Str. griseus*, *Str. rochei*, *Str. coelicolor* and *Str. Halstedii* according to the identification schemes of [21 and 17] and taking into consideration the scheme of Bergey’s Manual. Only 5 isolates were *Streptomyces* sp. due to low similarity indices (*Table 1*).

Members of the “other actinobacteria” group could be divided into 4 genera. The analysis of the first clusters (5 strains) showed morphological features such as well-developed, branched, septate mycelium with a diameter about 0.5 μm , and non-motile single spores, which were characteristic to the genus *Micromonospora* according to [8]. The members of this cluster showed an orange (young cells) to black-coloured (old cells) substrate mycelium phenotype, and no aerial mycelium was formed. After subse-

Table 1. Phenotypic and metabolic properties of NR-degrading actinobacteri.

Strain No.	Source	Species	NR	Hexadecane	Glucose	Fructose	Arabinose	Sucrose	Xylose
NR1	S	Str. griseus	+	-	+R	+	-	-	±
NR2	S	Str. griseus	+	-	+R	+	-	-	±
NR3	S	Str. griseus	+	-	+R	+	-	-	±
NR4	S	Str. griseus	+	-	+R	+	-	-	±
NR5	S	Str. griseus	+	-	+R	+	-	-	±
NR6	S	Str. griseus	+	-	+R	+	-	-	±
NR7	FW	Str. griseus	+	-	+R	+	-	-	±
NR8	FW	Str. griseus	+	-	+R	+	-	-	±
NR9	FW	Str. griseus	+	-	+R	+	-	-	±
NR10	FW	Str. griseus	+	-	+R	+	-	-	±
NR11	FWD	Str. griseus	+	-	+R	+	-	±	±
NR12	S	Str. rochei	+	-	+R	+	+	-	+
NR13	S	Str. rochei	+	-	+R	+	+	-	+
NR14	S	Str. rochei	+	-	+R	+	+	-	+
NR15	FW	Str. rochei	+	-	+R	+	+	-	+
NR16	FW	Str. rochei	+	-	+R	+	+	±	+
NR17	FWD	Str. rochei	+	-	+R	+	+	-	+
NR18	FWD	Str. rochei	+	-	+R	+	+	-	±
NR19	S	Str. coelicolor	+	-	+R	+	+	-	-
NR20	S	Str. coelicolor	+	-	+R	+	+	-	-
NR21	S	Str. coelicolor	+	-	+R	+	+	-	-
NR22	S	Str. coelicolor	+	-	+R	+	+	±	-
NR23	FW	Str. coelicolor	+	-	+R	+	+	-	-
NR24	FW	Str. halstedii	+	-	+R	+	+	±	-
NR25	FWD	Str. halstedii	+	-	+R	+	+	-	-
NR26	FWD	Str. halstedii	+	-	+R	+	+	-	±
NR27	S	Streptomyces sp.	+	-	+R	+	+	-	+
NR28	S	Streptomyces sp.	+	-	+R	+	+	-	+
NR29	S	Streptomyces sp.	+	-	+R	+	+	-	+
NR30	FW	Streptomyces sp.	+	-	+R	+	+	-	+
NR31	FWD	Streptomyces sp.	+	-	+R	+	+	-	+
NR32	S	Micromonospora aurantiaca	+	-	±R	-	+	-	-
NR33	FW	Micromonospora aurantiaca	+	-	±R	±	+	±	±
NR34	FWD	Micromonospora aurantiaca	+	-	+R	±	+	±	±
NR35	FWD	Micromonospora aurantiaca	+	-	+R	±	+	±	±
NR36	FWD	Micromonospora aurantiaca	+	-	+R	±	+	±	±
NR37	FW	Actinoplanes italicus	+	-	±R	-	+	±	±
NR38	FWD	Actinoplanes italicus	+	-	+R	+	+	+	±
NR39	FWD	Actinoplanes italicus	+	-	+R	+	+	+	±
NR40	S	Gordona sp.	+	-	+R	±	+	+	-
NR41	FW	Gordona sp.	+	-	+R	±	+	+	-
NR42	S	Nocardia sp.	+	-	+R	±	+	-	-

S: Soil, FW: Fresh water, FWD: Fresh water sediment +: good growth/halo formation

±: poor growth/halo formation

-: same growth as on mineral medium without carbon source/no halo

RF: spore chain rectusflexibilis

SP: spiral spore chain

R: repression of natural rubber degrading activity

Cont.

Strain No.	Source	Species	Inositol	Mannitol	Rhamnose	Raffinose	Aerial mycelium	Substrate mycelium	Spore chain
NR1	S	Str. griseus	-	+	-	-	yellow	light brown	RF
NR2	S	Str. griseus	-	+	-	-	yellow	light brown	RF
NR3	S	Str. griseus	-	+	-	-	yellow	light brown	RF
NR4	S	Str. griseus	-	+	-	-	yellow	light brown	RF
NR5	S	Str. griseus	-	+	-	-	yellow	light brown	RF
NR6	S	Str. griseus	-	+	-	-	yellow	light brown	RF
NR7	FW	Str. griseus	-	+	-	-	yellow	light brown	RF
NR8	FW	Str. griseus	-	+	-	-	yellow	light brown	RF
NR9	FW	Str. griseus	-	+	-	-	yellow	light brown	RF
NR10	FW	Str. griseus	-	+	-	-	grey	light brown	RF
NR11	FWD	Str. griseus	-	+	-	-	grey	light brown	RF
NR12	S	Str. rochei	+	+	+	-	grey	dark brown	SP
NR13	S	Str. rochei	+	+	+	-	grey	dark brown	SP
NR14	S	Str. rochei	+	+	+	-	grey	dark brown	SP
NR15	FW	Str. rochei	+	+	+	-	grey	dark brown	SP
NR16	FW	Str. rochei	+	+	+	-	grey	dark brown	SP
NR17	FWD	Str. rochei	+	+	+	-	grey	dark brown	SP
NR18	FWD	Str. rochei	+	+	-	-	grey	dark brown	SP
NR19	S	Str. coelicolor	-	+	-	-	yellow	dark brown	RF
NR20	S	Str. coelicolor	-	+	-	-	yellow	dark brown	RF
NR21	S	Str. coelicolor	-	+	-	-	yellow	dark brown	RF
NR22	S	Str. coelicolor	-	+	-	-	yellow	dark brown	RF
NR23	FW	Str. coelicolor	-	+	-	-	yellow	dark brown	RF
NR24	FW	Str. halstedii	+	+	-	-	grey	dark brown	RF
NR25	FWD	Str. halstedii	-	+	+	-	grey	dark brown	RF
NR26	FWD	Str. halstedii	-	+	+	-	grey	dark brown	SP
NR27	S	Streptomyces sp.	+	+	+	-	grey	light brown	SP
NR28	S	Streptomyces sp.	+	+	+	-	grey	light brown	SP
NR29	S	Streptomyces sp.	+	+	+	-	grey	light brown	SP
NR30	FW	Streptomyces sp.	+	+	+	-	grey	dark brown	RF
NR31	FWD	Streptomyces sp.	+	-	+	-	grey	dark brown	RF
NR32	S	Micromonospora aurantiaca	-	-	-	+	-	orange	mono-spore
NR33	FW	Micromonospora aurantiaca	-	-	-	+	-	orange	mono-spore
NR34	FWD	Micromonospora aurantiaca	-	-	-	+	-	orange-black	mono-spore
NR35	FWD	Micromonospora aurantiaca	-	-	-	+	-	orange-black	mono-spore
NR36	FWD	Micromonospora aurantiaca	-	-	-	+	-	orange-black	mono-spore
NR37	FW	Actinoplanes italicus	-	-	-	+	-	orange	mono-spore
NR38	FWD	Actinoplanes italicus	-	-	-	+	-	orange	mono-spore
NR39	FWD	Actinoplanes italicus	-	-	-	+	-	orange	mono-spore
NR40	S	Gordona sp.	-	-	-	+	-	orange to red	mono-spore
NR41	FW	Gordona sp.	-	-	-	+	-	orange to red	mono-spore
NR42	S	Nocardia sp.	-	-	-	+	-	orange	mono-spore

quent analysis of this cluster, it could be identified as *M. aurantiaca*. Koch et al. described this bacterium as a new species [9].

The second cluster was identified as *Actinoplanes* (3 isolates). Colonies had orange colour and produced pink to cherry coloured pigments diffusing into the medium. According to the Bergey's Manual of Systematic Bacteriology, this cluster is *Actinoplanes italicus*.

The third cluster was identified as *Gordona* (2 isolates). These isolates had rod shaped or coccoid cells. The genus *Gordona* is assigned to the suprageneric but phylogenetically coherent group of mycolic acid containing Actinobacteria [7]. These two isolates need more molecular analysis to reach the species level.

The last cluster (one isolate) had mycelia that eventually fragmented into rod shaped or coccoid cells. These characteristics class this isolate into the genus *Nocardia*. The present strain seems to be a newtype strain of the genus *Nocardia* and it needs more investigation.

This study demonstrates that certain bacteria can use the hydrocarbon of NR as a sole source of carbon and energy. These microorganisms may play an ecological role in the environment by mineralising NR latexes. Although no attempt was made to isolate the Actinobacteria selectively, all of the isolated rubber metabolising microorganisms were identified as *Streptomyces*, *Micromonospora*, *Gordona* and *Nocardia*. It suggests that degradation of NR is the privilege of mycelium forming microorganisms. Our results are consistent with those indicate that rubber degrading species of these genera are widely distributed in soils, water and fresh and marine sediments [5, 23 and 3]. Some of these isolates may have enough degrading potential that enables biotechnological use particularly in rubber products.

Regulation of NR degradation

The ability of NR-degrading bacteria to use low molecular mass monomers and high molecular mass polymer as carbon sources was tested (*Table 1*). When clearing zone formation was studied on NR plates containing one additional soluble carbon source, evidence for inhibition of NR degrading enzyme expression was obtained. Carbon sources that allow good growth e.g. glucose repressed the NR-degrading enzyme in most strains (*Table 1*). The extent of inhibition varied with strains and substrates. The biochemical mechanism of NR degradation has not been investigated. Since NR is a high molecular mass compound, which is too large to be taken by bacteria, as a first step the polymer has to be cleaved extracellularly. The extracellular nature of such enzyme system was shown by the appearance of translucent halos on latex containing solid media.

REFERENCES

- [1] Holt, J.G. et al., (eds.) (1989): *Bergey's Manual of Systematic Bacteriology – Williams and Wilkins, Baltimore.*
- [2] Cundell, A.M. and Mulcock, A. P. (1973): The effect of curing agent concentration on the microbiological deterioration of vulcanized natural rubber. – *International Biodeterioration Bulletin* 9: 91-94.
- [3] Hanstveit, A. O., Gerritse, G. A. and Scheffers, W. A. (1986): A study of the biodegradation of vulcanized rubber sealing, exposed to inoculated tap water. In: L. H.

- G. Morton (ed.), The biodeterioration of constructional material. Proceeding of the Biodeterioration Society Meeting, 18-19 September 1986, Delft, The Netherlands, Biodeterioration Society, Slough, United Kingdom. p. 87-96.
- [4] Heisey, R. M. and Papadatos, S. (1995): Isolation of microorganisms able to metabolise purified natural rubber. – *Applied Environmental Microbiology* 61: 3092-3097.
- [5] Hutchinson, M., Ridgway, J. W and Cross, T. (1975): Biodeterioration of rubber in contact with water, sewage and soil, p.187–202. In: R.J. Gilbert and D.W. Lovelock (ed.). *Microbial aspects of the deterioration of materials*. Academic press, New York.
- [6] Jang, J.W., Yoo, T. S., Oh, J. H. and Iwasaki, I. (1998): Discarded tire recycling practices in the United States, Japan and Korea. – *Research Conservation Recycle* 22: 1-14.
- [7] Klalte, S., Rainey, F. A., and Kroppenstedt, R. M. (1994): Transfer of *Rhodococcus aichiensis* Tsukamura 1982 and *Nocardia amarae* Lechevalier and Lechevalier 1974 to the genus *Gordana* as *Gordana aichiensis* comb. Nov. and *Gordana amarae* comb. Nov. – *International Journal Systematic Bacteriology* 44: 769-773.
- [8] Kawamoto, I. (1989): Genus *Micromonospora* Orskov 1932, p. 2442-2450. In: T. Williams, M. E. Sharpe, and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 4. The Williams and Wilkins Co., Baltimore.
- [9] Koch, C., Kroppenstedt, R. M and Stackebrandt, E. (1996): Intrageneric relationships of the actinomycete genus *Micromonospora*. – *International Journal Systematic Bacteriology* 36: 383-387.
- [10] Leeflang, K.W.H (1963): Microbiologic degradation of rubber. – *Journal American Water Works Association* 53: 1523-1535.
- [11] Linos, A., Reichelt, R., Keller, U. and Steinbuchel, A. (2000): A Gram-negative bacterium, identified as *Pseudomonas aeruginosa* AL 98, is a potent degrader of natural rubber and synthetic cis-1,4-polyisoprene. – *FEMS Microbiology. Letters* 182: 155-161.
- [12] Linos, A., Steinbuchel, A., Sproer, C. and Kroppenstedt, R. M. (1999): *Gordana polyisoprenivorans* sp. nov. A rubber degrading actinomycete isolated from automobile tire. – *International Journal Systematic Bacteriology* 49: 1785-1791.
- [13] Liu, H. S., Mead, J. L. and Stacer, R. G. (2000): Environmental effects of recycled rubber in land-fill applications.- *Rubber Chemistry Technology* 73: 551-564.
- [14] Smith, F. G and Klingensmith, W. B (1991): Tires as a consumer and source of recycled material. – *Rubber chemistry Technology* 64: 122-130.
- [15] Sohngen, N. L., and Fol, J. G. (1914): Die zersetzung des kautschuks darc/mikroben. – *Zentral Bakteriologie Parasitenkd Infektionskr Hygiene* 40: 87-98.
- [16] Sokal, R.R., Michener, C.D. 1958. A statistical method for evaluating systematic relationships. *Univ. Kans. Sci. Bull.* 38, 1409-1438.
- [17] Szabo, I. M., Marton, M., and Buti, I. C. (1978): A diagnostic key for the identification of species of *Streptomyces* and *Streptoverticillium* included in the International *Streptomyces* Projects. – *A Bot. Academic Science Hungary* 21: 387-418.
- [18] Tsuchii, A. (1995): Microbial degradation of natural rubber. In: *Biotransformations: Microbial degradation of health risk compounds* (Singh, Ved

- Pal, Ed.). – Progress Indian Microbiology 32: 177-187.
- [19] Tsuchii, A. and Takeda, K. (1990): Rubber-degrading enzyme from a bacteria culture. – Applied Environmental Microbiology 56: 269-274.
- [20] Tsuchii, A. and Tokiwa, Y. (2001): Microbial degradation of tyre rubber particles.- Biotechnology Letters 23: 963-969.
- [21] Williams, S. T. Goodfellow, N., Aldreson, G. Wellington, E. M. H., Sneath, P. H. A., and Sackin, M. J. (1983): Numerical classification of Streptomyces and related genera. – Journal General Microbiology 129: 1743-1813.
- [22] Zobell, C.E. and Beckwith, J. D. (1994): The deterioration of rubber products by microorganisms. – Journal American Water Works Association 36: 439-453.
- [23] Zyska, B. J. (1981): Rubber, pp. 323-385. In A. H. Rose (ed.), Microbial biodeterioration, economic microbiology, vol. 6. Academic Press, London, England.